



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 9/127	A1	(11) International Publication Number: WO 95/13796 (43) International Publication Date: 26 May 1995 (26.05.95)
(21) International Application Number: PCT/US94/12957 (22) International Filing Date: 10 November 1994 (10.11.94) (30) Priority Data: 08/153,657 16 November 1993 (16.11.93) US (71) Applicant: DEPOTECH CORPORATION [US/US]; 11025 North Torrey Pines Road, La Jolla, CA 92037 (US). (72) Inventors: SANKARAM, Mantripragada; Apartment #53, 12726 Torrey Bluff Drive, San Diego, CA 92130 (US). KIM, Sinil; 548 Ford Avenue, Solana Beach, CA 92705 (US). (74) Agents: BOSTICH, June, M. et al.; Spensley Horn Jubas & Lubitz, 5th floor, 1880 Century Park East, Los Angeles, CA 90067 (US).		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ). Published <i>With international search report.</i>
(54) Title: VESICLES WITH CONTROLLED RELEASE OF ACTIVES (57) Abstract A synthetic membrane vesicle composition containing at least one release-rate modifying agent other than a hydrohalide and at least one biologically active substance, the vesicles having defined size distribution, adjustable average size, internal chamber size and number, and a controlled release rate of the biologically active substance. A process for making the composition features addition of a release-rate modifying agent effective to prolong, sustain, and control the rate of release from the vesicles of the biologically active substance at therapeutic levels after encapsulation.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

VESICLES WITH CONTROLLED RELEASE OF ACTIVES

BACKGROUND OF THE INVENTION

5 **1. Field of the Invention**

The invention relates to compositions of synthetic membrane vesicles useful as a drug delivery system and to processes for their manufacture.

2. Background of the Invention

10 Multivesicular liposomes are one of the three main types of liposomes, first made by Kim, et al. (*Biochim. Biophys. Acta*, 782:339-348, 1983), and are uniquely different from other lipid-based drug delivery systems such as unilamellar (Huang, *Biochemistry*, 8:334-352, 15 1969; Kim, et al., *Biochim. Biophys. Acta*, 646:1-10, 1981) and multilamellar (Bangham, et al., *J. Mol. Bio.*, 13:238-252, 1965) liposomes. In contrast to unilamellar liposomes, multivesicular particles contain multiple aqueous chambers per particle. In contrast to 20 multilamellar liposomes, the multiple aqueous chambers in multivesicular particles are non-concentric.

The prior art describes a number of techniques for producing unilamellar and multilamellar liposomes; for example, U.S. Patent No. 4,522,803 to Lenk; 4,310,506 25 to Baldeschwieler; 4,235,871 to Papahadjopoulos; 4,224,179 to Schneider; 4,078,052 to Papahadjopoulos; 4,394,372 to Taylor; 4,308,166 to Marchetti; 4,485,054 to Mezei; and 4,508,703 to Redziniak. The prior art also describes methods for producing multivesicular 30 liposomes that proved unstable in biological fluids (Kim, et al., *Biochim. Biophys. Acta*, 728:339-348, 1983). For a comprehensive review of various methods of unilamellar and multilamellar liposome preparation, refer to Szoka, et al., *Ann. Rev. Biophys.* 35 *Bioeng.*, 9:465-508, 1980.

In the method of Kim, et al. (*Biochim. Biophys. Acta*, 728:339-348, 1983), the encapsulation efficiency

- 2 -

of small molecules, such as cytosine arabinoside, was low, and had rapid release rate in biological fluids. Subsequent studies (Kim, et al., *Cancer Treat. Rep.*, 71:705-711, 1987) showed that the rapid release rate of encapsulated molecules in biological fluids can be improved by encapsulating in the presence of a hydrochloride.

Optimal treatment with many drugs requires maintenance of a drug level for a prolonged period of time. For example, optimal anti-cancer treatment with cell cycle-specific antimetabolites requires maintenance of a cytotoxic drug level for a prolonged period of time. Cytarabine is a highly scheduled-dependent anti-cancer drug. Because this drug kills cells only when they are replicating DNA, a prolonged exposure at therapeutic concentration of the drug is required for optimal cell kill. Unfortunately, the half-life of Cytarabine after an intravenous (IV) or subcutaneous (SC) dose is very short, with the half-life in the range of a few hours. To achieve optimal cancer cell kill with a cell cycle phase-specific drug like Cytarabine, two major requirements need to be met: first, the cancer must be exposed to a high concentration of the drug without doing irreversible harm to the host; and second, the tumor must be exposed for a prolonged period of time so that all or most of the cancer cells have attempted to synthesize DNA in the presence of Cytarabine.

Heretofore, control of the release rate was inflexible, and the choice of release-rate modifying agents was limited primarily to hydrohalides. For a drug-delivery system, it is highly advantageous to be flexible in controlling the release rate for encapsulated substances and to have a wide choice of release-rate modifying agents.

Accordingly, it is an object of the present invention to provide a slow-releasing depot preparation

- 3 -

which provides a prolonged and sustained exposure of a biologically active substance at a therapeutic concentration, with a controlled release rate.

It is a further object of the present invention to provide a method of preparing such depot preparations.

Other and further objects, features, and advantages of the invention are inherent therein and appear throughout the specification and claims.

10 SUMMARY OF THE INVENTION

The compositions of the present invention comprise synthetic membrane vesicles, i.e. lipid vesicles with multiple internal aqueous chambers formed by non-concentric layers and wherein the chambers contain one or more release-rate modifying agents effective in slowing the release rate of the encapsulated biologically active substances. The present invention also provides methods of making such compositions.

The present synthetic membrane vesicle compositions have high encapsulation efficiency, controlled release rate of the encapsulated substance, well defined, reproducible size distribution, spherical shape, adjustable average size that can be easily increased or decreased, adjustable internal chamber size and number.

The process for producing these compositions comprises (1) mixing one or more volatile organic solvents and a lipid component containing at least one neutral lipid and at least one amphipathic lipid having one or more net negative charges; (2) adding into the organic solvent an immiscible first aqueous component containing one or more biologically active substances to be encapsulated; (3) adding to either or both the organic solvent and the first aqueous component, a release-rate modifying agent effective in slowing the release rate of the encapsulated biologically active substances; (4) forming a water-in-oil emulsion from the two immiscible components; (5) immersing the water-

- 4 -

in-oil emulsion into a second immiscible aqueous component; (6) dispersing the water-in-oil emulsion to form solvent spherules containing in them multiple droplets of the first aqueous component; and (7) removing the organic solvents, such as by evaporation, from the solvent spherules to form the synthetic membrane vesicles. Addition of one or more release-rate modifying agents effective in slowing the release rate of the encapsulated biologically active substances in biological fluids and *in vivo* is essential.

A BRIEF DESCRIPTION OF THE DRAWING

Figure 1 is a graph showing the rate of release of a drug from synthetic membrane vesicles suspended in human plasma at 37°C. The symbols used indicate the release rate modifying agent employed and are identified in Table 2.

DESCRIPTION OF THE PREFERRED EMBODIMENT

The term "synthetic membrane vesicles" as used throughout the specification and claims means man-made, microscopic lipid-vesicles consisting of lipid bilayer membranes, enclosing multiple non-concentric aqueous chambers. In contrast, unilamellar liposomes have a single aqueous chamber; and multilamellar liposomes have multiple "onion-skin" type of concentric membranes, in between which are shell-like concentric aqueous compartments.

The term "solvent spherule" as used throughout the specification and claims means a microscopic spheroid droplet of organic solvent, within which is multiple smaller droplets of aqueous solution. The solvent spherules are suspended and totally immersed in a second aqueous solution.

The term "neutral lipid" means oil or fats that have no membrane-forming capability by themselves and lack a hydrophilic "head" group.

- 5 -

The term "amphipathic lipids" means those molecules that have a hydrophilic "head" group and hydrophobic "tail" group and have membrane-forming capability.

5 The term "release-rate modifying agent" means molecules other than hydrohalides added during the process of making or manufacturing the synthetic membrane vesicles that are effective in either slowing or increasing the release rate of the encapsulated biologically active substances from the synthetic
10 membrane vesicles.

Briefly, the method of the invention comprises making a "water-in-oil" emulsion by (1) dissolving amphipathic lipids in one or more volatile organic solvents for the lipid component, (2) adding to the
15 lipid component an immiscible first aqueous component and a biologically active substance to be encapsulated, and (3) adding to either or both the organic solvent and the first aqueous component, a release-rate modifying agent effective in slowing the release rate
20 of the encapsulated biologically active substances from the synthetic membrane vesicles, and then emulsifying the mixture mechanically.

In the emulsion, the water droplets suspended in the organic solvent will form the internal aqueous
25 chambers, and the monolayer of amphipathic lipids lining the aqueous chambers will become one leaflet of the bilayer membrane in the final product. The emulsion is then immersed in a second aqueous component containing one or more nonionic osmotic agents and an
30 acid-neutralizing agent of low ionic strength, such as a proton acceptor preferably selected from free-base lysine, free-base histidine, or a combination thereof. Then the emulsion is agitated either mechanically, by ultrasonic energy, nozzle atomizations, and the like,
35 or by combinations thereof, to form solvent spherules suspended in the second aqueous component.

- 6 -

The solvent spherules contain multiple aqueous droplets with the substance to be encapsulated dissolved in them. The organic solvent is removed from the spherules, preferably by evaporation of a volatile solvent, for instance by passing a stream of gas over the suspension. When the solvent is completely removed, the spherules convert into synthetic membrane vesicles. Representative gases satisfactory for use in evaporating the solvent include nitrogen, helium, argon, oxygen, hydrogen, and carbon dioxide.

The release-rate modifying agent is any molecule that is effective in slowing the rate of release of the encapsulated biologically active substances from the synthetic membrane vesicles in biological fluids and *in vivo*, with the result that the release rate of the substances is slower than that from synthetic membrane vesicles produced in the absence of such a release-rate modifying agent. The release-rate modifying agents include, but are not limited to, perchloric acid, nitric acid, formic acid, acetic acid, trifluoroacetic acid, trichloroacetic acid, sulfuric acid, phosphoric acid, and combinations thereof. The amounts of the release-rate modifying agents used is one effective to provide a prolonged, sustained, and controlled rate of release at therapeutic levels of the encapsulated biologically active substances. For example, the concentration of the release-rate modifying agent in the the organic solvent or the first aqueous component to which it is added is in the range from about 0.1 mM to about 0.5 M and preferably from about 10 mM to about 200 mM.

Many different types of volatile hydrophobic solvents such as ethers, hydrocarbons, halogenated hydrocarbons, or Freons may be used as the lipid-phase solvent. For example, diethyl ether, isopropyl and other ethers, chloroform, tetrahydrofuran, halogenated

- 7 -

ethers, esters and combinations thereof are satisfactory.

5 In order to prevent the solvent spherules from sticking to each other and to the vessel wall, it is preferred that at least 1 percent molar ratio of an amphipathic lipid with a net negative charge be included in the spherules, that the suspending second aqueous solution have a very low ionic strength, and, when an acid is used, that an agent for neutralizing
10 the acid be added to the second aqueous solution to form a concentration of from about 0.1 mM to about 0.5 M therein to prevent coalescence of the solvent spherules to form a messy scum. In addition, one or more nonionic osmotic agents, such as trehalose,
15 glucose, or sucrose, may optionally be used in the suspending aqueous solution to keep the osmotic pressure within and without the membrane vesicles balanced.

Various types of lipids can be used to make the
20 synthetic membrane vesicles, and the only two requirements are that one amphipathic lipid with a net negative charge and a neutral lipid be included. Examples of neutral lipids are triolein, trioctanoin, vegetable oil such as soybean oil, lard, beef fat,
25 tocopherol, and combinations thereof. Examples of amphipathic lipids with net negative charge are cardiolipin, the phosphatidylserines, phosphatidylglycerols, and phosphatidic acids.

The second aqueous component is an aqueous solution
30 containing low ionic strength solutes such as carbohydrates including glucose, sucrose, lactose, and amino acids such as lysine, free-base histidine and combinations thereof.

35 Many and varied biological substances and therapeutic agents can be incorporated by encapsulation within the synthetic membrane vesicles.

- 8 -

The term "therapeutic agent" as used herein for the compositions of the invention includes, without limitation, drugs, radioisotopes, and immunomodulators. Similar substances are known or can be readily
5 ascertained by one of skill in the art. There may be certain combinations of therapeutic agent with a given type of synthetic membrane vesicles that are more compatible than others. For example, the method for producing the synthetic membrane vesicles may not be
10 compatible with the continued biological activity of a proteinaceous therapeutic agent. However, since conditions that would produce an incompatible pairing of a particular therapeutic agent with a particular dispersion system are well known, or easily
15 ascertained, it is a matter of routine to avoid such potential problems.

The drugs that can be incorporated into the dispersion system as therapeutic agents include non-proteinaceous as well as proteinaceous drugs. The term
20 "non-proteinaceous drugs" encompasses compounds that are classically referred to as drugs, such as mitomycin C, daunorubicin, vinblastine, AZT, and hormones. Of particular interest are anti-tumor cell-cycle specific drugs such as cytarabine, methotrexate, 5-fluorouracil
25 (5-FU), floxuridine (FUDR), bleomycin, 6-mercapto-purine, 6-thioguanine, fludarabine phosphate, vincristine, and vinblastine.

Examples of proteinaceous materials that can be incorporated into the synthetic membrane vesicles are
30 DNA, RNA, proteins of various types, protein hormones produced by recombinant DNA technology effective in humans, hematopoietic growth factors, monokines, lymphokines, tumor necrosis factor, inhibin, tumor growth factor alpha and beta, Mullerian inhibitory
35 substance, nerve growth factor, fibroblast growth factor, platelet-derived growth factor, pituitary and hypophyseal hormones including LH and other releasing

- 9 -

hormones, calcitonin, proteins that serve as immunogens for vaccination, and DNA and RNA sequences.

The following TABLE 1 includes a list of representative biologically active substances effective in humans that can be encapsulated in synthetic membrane vesicles in the presence of a release-rate modifying agent of the invention, and also includes biologically active substances effective for agricultural uses.

10

TABLE 1

	<u>Antiasthmas</u>	<u>Antiarrhythmics</u>	<u>Tranquilizers</u>
	metaproterenol	propanolol	chlorpromazine
15	aminophylline	atenolol	benzodiazepine
	theophylline	verapamil	butyrophenones
	terbutaline		hydroxyzines
	norepinephrine	<u>Antianginas</u>	meprobamate
	ephedrine	isosorbide dinitrate	phenothiazines
20	isoproterenol		thioxanthenes
	adrenalin		
	<u>Cardiac glycosides</u>	<u>Hormones</u>	<u>Steroids</u>
	digitalis	thyroxine	prednisone
25	digitoxin	corticosteroids	triamcinolone
	lanatoside C	testosterone	hydrocortisone
	digoxin	estrogen	dexamethasone
		progesterone	betamethasone
30		mineralocorticoid	prednisolone
	<u>Antihypertensives</u>	<u>Antidiabetics</u>	<u>Antihistamines</u>
	apresoline	Diabenese	pyribenzamine
	atenolol	insulin	
	chlorpheniramine		
35	captopril		diphenhydramine
	reserpine		
	<u>Antiparasitics</u>	<u>Antineoplastics</u>	<u>Sedatives and</u>
	<u>Analgesics</u>		
40	praziquantel	azathioprine	morphine
	metronidazole	bleomycin	dilaudid
	pentamidine	cyclophosphamide	codeine
	ivermectin	vincristine	codeine-like
	synthetics		
45	<u>Nucleic Acids and Analogs</u>	methotrexate	demerol
	DNA	6-TG	oxymorphone
	RNA	6-MP	phenobarbital
		vinblastine	barbiturates

- 10 -

	methyolphosphonates and analogs	VP-16	fentanyl
	Antisense nucleic acids	VM-26	ketorolac
5		cisplatin	
		5-FU	
		FUDR	
		fludarabine phosphate	
	<u>Antibiotics</u>	<u>Immunomodulators</u>	<u>Vasopressors</u>
10	penicillin	interferon	dopamine
	tetracycline	interleukin-2	dextroamphetamine
	amikacin	gammaglobulin	
	erythromycin	monoclonal antibodies	
	cephalothin		
	imipenem	<u>Antifungals</u>	<u>Antivirals</u>
15	cefotaxime	amphotericin B	acyclovir and derivatives
	carbenicillin	myconazole	Gancyclovir and phosphates
	ceftazidime	muramyl dipeptide	Winthrop-51711
	kanamycin	clotrimazole	ribavirin
	tobramycin	ketoconazole	rimantadine/amantadine
20	ampicillin	fluconazole	azidothymidine & derivates
	gentamycin	itraconazole	adenine arabinoside
	cefoxitin		amidine-type protease
	cefadroxil		inhibitors
	cefazolin		
25	other aminoglycosides		
	amoxicillin		
	moxalactam		
	piperacillin		
	vancomycin		
30	ciprofloxacin		
	other quinolones		
	<u>Vaccines</u>		
35	other recombinant, killed and live vaccines and antigenic material for use as vaccines.		
	antigenic material for the treatment of allergies		
	influenza		
	respiratory syncytial virus		
	HIV vaccine		
40	Hemophilus influenza vaccines		
	Hepatitis A,B,C vaccines		
	mumps		
	rubella		
	measles		
45	tetanus		
	malaria vaccines		
	herpes		
	cancer vaccines		
	Anti-leu-3a vaccine		
50			
	<u>Monoclonal Antibodies</u> (human, mouse other species-derived and/or recombinant and/ or fusions and/or fragments thereof)		
	OKT3		
	OKT4		
55	HA-1A		
	Anti-Carcino-Embryonic Antigen Antibodies		

- 11 -

- Anti-Ganglioside Antibodies: Anti GD2, Anti GM2, Anti GD3, Anti GM3
 Urinary Tract-Associated Antigen-related antibodies
 Anti-IL-2 Receptor
 Chimeric Anti-Leu-2
 5 Anti-IL-2 receptor
 Anti-Leu-2
 Chimeric Anti-Leu-3a
 Chimeric L6
 MAb-L6
 10 Radiolabeled L6
 Centorex
 Centoxin
 Panorex
 Anti-LPS
 15 Immunotoxin
 Anti-tumor necrosis factor
 Anti-pseudomonas
 Anti-tumor necrosis factor
 OncoRad 103
 20 OncoScint CR103
 OncoScint OV103
 OncoScint PR356
 OncoTher 130
 KS 1/4-DAVLB
 25 ADCC agent
 Murine monoclonal antibodies to human B-cell lymphomas (anti-
 idiotypes)
 Murine monoclonal antibody (lMelpgl) (anti-idiotypic) against murine
 monoclonal antibody to melanoma-associated antigen
 30 Anti-B4-blocked ricin
 Anti-My9-blocked ricin
 ImmuRaid-CEA
 MAb against colorectal, ovarian, and lung cancers
 rhenium-186 MAb
 35 Orthoclone OKT®
 E5™
 LYM-1
 TNT
 XomaZyme®-791
 40 XomaZyme®-CD5 Plus
 XomaZyme®-CD7 Plus
 XomaZyme®-Mel
- Herbicides
 45 Triazine
 chloroacetamide
 cyanazine
 bentazone
 Roundup
 50 Rodeo
 butachlor
 CNP
 chlomethoxynil
 simetryne
 55 Atrazine
 Alachlor

- 12 -

Cyanazine
metolachlor
metribuzin
5 phenoxy herbicides: 2,4-D [(2,4-dichlorophenoxy)acetic acid],
 2,4-D amine (2,4-dichlorophenoxyacetic acid dimethylamine),
 2,4-D isooctyl (2,4-dichlorophenoxyacetic acid isooctyl ester),
 2,4,5-T amine (2,4,5-trichlorophenoxyacetic acid trimethylamine)
other triazine herbicides
other chloroacetamide herbicides
10 other phenoxyacid herbicides

Pesticides

Abamectin
other avermectins
15 atrazine
 lindane
 dichlorvos
 dimethoate
 warfarin
20 p,p'-DDD
 p,p'-DDE
 HCH
 DMDT
 aldrin
25 dieldrin
 Aldicarb
 EDB
 DCP
 DBCP
30 simazine
 cyanazine
 Bacillus thuringiensis toxin
 Bacillus thuringiensis var. kurstaki
 bis(tri-n-butyltin)oxide (TBTO)
35 other organochlorine pesticides

Proteins and Glycoproteins

lymphokines
 interleukins - 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11.
40 cytokines
 GM-CSF
 M-CSF
 G-CSF
tumor necrosis factor
45 inhibin
 tumor growth factor
 Mullerian inhibitors substance
 nerve growth factor
 fibroblast growth factor
50 platelet derived growth factor
 coagulation factors (e.g. VIII, IX, VII)
 insulin
 tissue plasminogen activator
 histocompatibility antigen
55 oncogene products
 myelin basic protein

- 13 -

collagen
fibronectin
laminin
other proteins made by recombinant DNA
5 technology
erythropoietin
IL-3/GM-CSF fusion proteins
Monoclonal antibodies
Polyclonal antibodies
10 antibody-toxin fusion proteins
antibody radionuclide conjugate
Interferons
Fragments and peptide analogs, and analogs of fragment of proteins,
peptides
15 and glycoproteins.
Epidermal growth factor
CD4 receptor and other recombinant receptors
other proteins isolated from nature
Antidiuretic hormone
20 oxytocin
adrenocorticotropin Hormone
calcitonin
follicle stimulating hormone
luteinizing hormone releasing hormone
25 luteinizing hormone
gonadotrophin
transforming growth factors
streptokinase
Human Growth Hormone,
30 Somatotropins for other species, including, but not limited to:
 1. Porcine,
 2. Bovine,
 3. Chicken,
 4. Sheep,
35 5. Fish,
Growth Hormone releasing hormones for humans and various animal
species,
Glucagon,
Desmopressin,
40 Thyroid Releasing Hormone,
Thyroid Hormone,
Secretin,
Magainins,
Integrins,
45 Adhesion Peptides, including, but not limited to, those having the
 Arginine-Glutamine-Aspartic Acid sequence,
Super Oxide Dismutase,
Defensins,
T-Cell Receptors,
50 Bradykinin antagonists,
Pentigetide,
Peptide T,
Antinflammins,
Major Histocompatibility (MHC) complex components and peptides
55 targeted to the MHC,
Protease inhibitors,

- 14 -

- 5 Lypressin,
 Buserelin,
 Leuprolide,
 Nafarelin,
 10 Deslorelin,
 Goserelin,
 Historelin,
 Triptorelin,
 LHRH antagonists,
 15 HOE-2013,
 Detirelix,
 Org-30850,
 ORF-21243,
 Angiotensin Converting Enzyme inhibitor Peptide,
 20 Renin inhibitory peptides,
 Ebiratide (HOE-427),
 DGAVP,
 Opiate receptor agonists and antagonists, including, but not
 limited to:
 1. Enkephalins,
 2. Endorphins,
 E-2078,
 DPDPE,
 Vasoactive intestinal peptide,
 25 Atrial Natriuretic Peptide,
 Brain Natriuretic Peptide,
 Atrial Peptide clearance inhibitors,
 Hirudin,
 Oncogene Inhibitors,
 30 Other Colony Stimulating Factors,
- | | <u>Neurotransmitters</u> | <u>Radionuclides</u> | <u>Radio contrasts</u> |
|----|--------------------------------|----------------------|------------------------|
| | Dopamine | Technetium | Gadolinium chelates |
| | Epinephrine | Indium | Iohexol |
| 35 | Norepinephrine | Yttrium | Ethiodol |
| | acetylcholine | Gallium | Iodexinol |
| | Gammaamino butyric acid | | |
| | <u>Others</u> | | |
| 40 | cell surface receptor blockers | | |

The term "therapeutically effective" as it pertains
 to the compositions of the invention means that a
 45 therapeutic agent is present in the aqueous phase
 within the vesicles at a concentration sufficient to
 achieve a particular medical effect for which the
 therapeutic agent is intended. Examples, without
 limitation, of desirable medical effects that can be
 50 attained are chemotherapy, antibiotic therapy, and
 regulation of metabolism. Exact dosages will vary

- 15 -

depending upon such factors as the particular therapeutic agent and desirable medical effect, as well as patient factors such as age, sex, general condition, and the like. Those of skill in the art can readily
5 take these factors into account and use them to establish effective therapeutic concentrations without resort to undue experimentation.

Generally, however, the dosage range appropriate for human use includes the range of 0.1-6000 mg/sq m of
10 body surface area. For some applications, such as subcutaneous administration, the dose required may be quite small, but for other applications, such as intraperitoneal administration, the dose desired to be used may be very large. While doses outside the
15 foregoing dose range may be given, this range encompasses the breadth of use for practically all the biologically active substances.

The synthetic membrane vesicles may be administered for therapeutic applications by any desired route; for
20 example, intramuscular, intrathecal, intraperitoneal, subcutaneous, intravenous, intralymphatic, oral and submucosal, under many different kinds of epithelia including the bronchial epithelia, the gastrointestinal epithelia, the urogenital epithelia,
25 and various mucous membranes of the body.

In addition the synthetic membrane vesicles of the invention can be used to encapsulate compounds useful in agricultural applications, such as fertilizers, pesticides, and the like. For use in agriculture, the
30 synthetic membrane vesicles can be sprayed or spread onto an area of soil where plants will grow and the agriculturally effective compound contained in the vesicles will be released by contact with rain and irrigation waters. Alternatively the slow-releasing
35 vesicles can be mixed into irrigation waters to be applied to plants and crops. One skilled in the art will be able to select an effective amount of the

- 16 -

compound useful in agricultural applications to accomplish the particular goal desired, such as the killing of pests, the nurture of plants, etc.

5 The synthetic membrane vesicles may be modified in order to impart organ or cell target specificity, for instance by incorporating them into a targeted delivery system. Such modifications may be particularly relevant for using the synthetic membrane vesicles of the invention to administer drugs that are highly toxic
10 or capable of inducing severe side effects, such as taxol.

 The targeting of the synthetic membrane vesicles is classified based on anatomical and mechanistic factors.
15 In anatomical targeting, the synthetic membrane vesicle is targeted to a specific body location, for example, organ-specific, cell-specific, and organelle-specific targeting. Mechanistic targeting can be distinguished based upon whether it is passive or active. Passive
20 targeting utilizes the natural tendency of the synthetic membrane vesicles of the invention to distribute to cells of the reticulo-endothelial system (RES) in organs which contain sinusoidal capillaries. In active targeting, on the other hand, the synthetic
25 membrane vesicle is incorporated into a targeted delivery system by coupling it to a specific ligand, such as a monoclonal antibody, sugar, glycolipid, or protein, or by changing the composition or size of the synthetic membrane vesicles in order to achieve
30 targeting to organs and cell types other than the naturally occurring sites of localization (see, for example, *Remington's Pharmaceutical Sciences*, Gannaro, A.R., ed., Mack Publishing, 18 Edition, pp. 1691-1693, 1990)

35 In general, the compounds to be bound to the surface of the synthetic membrane vesicles will be ligands and receptors that allow the dispersion system

- 17 -

to actively "home in" on the desired tissue. A ligand may be any compound of interest that will specifically bind to another compound, referred to as a receptor, such that the ligand and receptor form a homologous pair.

The surface of the targeted delivery system can be modified in a variety of ways. For instance, lipid groups can be incorporated into the lipid bilayer of the synthetic membrane vesicles in order to maintain the targeting ligand in stable association with the lipid bilayer. Various linking groups can be used for joining the lipid chains to the targeting ligand (Mannino, et al., *Bio Techniques*, 6(7):682, 1988). The compounds bound to the surface of the synthetic membrane vesicles may vary from small haptens of from about 125-200 molecular weight to much larger antigens with molecular weights of at least about 6000, but generally of less than 1 million molecular weight. Proteinaceous ligand and receptors are of particular interest.

In general, the surface membrane proteins that bind to specific effector molecules are referred to as receptors. In the present invention, the preferred receptors are antibodies. These antibodies may be monoclonal or polyclonal and may be fragments thereof such as Fab F(ab')₂, and F_v, which are capable of binding to an epitopic determinant. Techniques for binding of proteins, such as antibodies, to synthetic membrane vesicles are well known (see, for example, U.S. 4,806,466 and U.S. 4,857,735, incorporated by reference).

Antibodies can be used to target the synthetic membrane vesicles to specific cell-surface ligands. For example, certain antigens expressed specifically on tumor cells, referred to as tumor-associated antigens (TAAs) may be exploited for the purpose of targeting antibody-containing synthetic membrane vesicles

- 18 -

directly to malignant tumors. Since the composition incorporated into the synthetic membrane vesicles may be indiscriminate with respect to cell type in its action, a targeted synthetic membrane vesicles offers a significant improvement over randomly injecting non-specific synthetic membrane vesicles. A number of procedures can be used to covalently attach either polyclonal or monoclonal antibodies to a bilayer of the synthetic membrane vesicles. Antibody-targeted synthetic membrane vesicles can include monoclonal or polyclonal antibodies or fragments thereof such as Fab, or F(ab')₂, as long as they bind efficiently to the antigenic epitope on the target cells. Synthetic membrane vesicles may also be targeted to cells expressing receptors for hormones or other serum factors (Malone, et al., *Proc. Nat'l. Acad. Sci, USA*, 86:6077, 1989; Gregoriadis, *Immunology Today*, 11(3):89, 1990; both incorporated by reference).

The following examples illustrate the manner in which the invention can be practiced. It is understood, however, that the examples are for the purpose of illustration and the invention is not to be regarded as limited to any of the specific materials or conditions therein.

EXAMPLE 1

Step 1) In a clean glass cylinder (2.5 cm inner diameter X 10.0 cm height), 5 ml of a solution containing 46.5 μ moles of dioleoyl phosphatidylcholine, 10.5 μ moles of dipalmitoyl phosphatidylglycerol, 75 μ moles of cholesterol, 9.0 μ moles of triolein in chloroform were placed (the lipid phase).

Step 2) Five ml of aqueous phase, cytarabine (20 mg/ml) dissolved in 0.136 N perchloric acid, a release-rate modifying agent, is added into the above glass cylinder containing lipid phase. The osmolarity of the aqueous solution is about 274 ± 20 mOs/kg. For the

- 19 -

other release-rate modifying agents namely, nitric acid, formic acid, sulfuric acid, phosphoric acid, acetic acid, trichloroacetic acid, and trifluoroacetic acid, 20 mg/ml solutions of cytarabine were prepared with these agents to yield aqueous solutions that are nearly isotonic with respect to the final storage medium, namely normal saline (0.9% sodium chloride).

Step 3) For making the water-in-oil emulsion, a homogenizer (AutoHomoMixer, Model M, Tokushu Kika, Osaka, Japan) was used by mixing for 8 minutes at a speed of 9000 rpm.

Step 4) For making the chloroform spherules suspended in water, 20 ml of a solution containing 4 percent dextrose and 40 mM lysine was layered on top of the water-in-oil emulsion, and then mixed for 60 seconds at a speed of 4000 rpm to form the chloroform spherules.

Step 5) The chloroform spherule suspension in the glass cylinder was poured into the bottom of a 1000 ml Erlenmeyer flask containing 30 ml of water, glucose (3.5 g/100 ml), and free-base lysine (40 mM). A stream of nitrogen gas at 7 l/minute was flushed through the flask to slowly evaporate chloroform over 20 minutes at 37°C. 60 ml of normal saline (0.9% sodium chloride) was added to the flask. The synthetic membrane vesicles were then isolated by centrifugation at 600 X g for 10 minutes. The supernatant was decanted, and the pellet was resuspended in 50 ml of normal saline. The pellet was resuspended in saline to yield a final concentration of 10 mg Cytarabine per ml of suspension.

The average length-weighted mean diameter of the resulting synthetic membrane vesicles particles is in the range from 12-16 μm . Percentage of capture of Cytarabine is given in TABLE 2. The use of different release-modifying agents had marked influence on the rate of Cytarabine release from the synthetic membrane vesicles incubated in human plasma. The percent of

- 20 -

Cytarabine retained in the synthetic membrane vesicles after incubation at 37°C in human plasma for the different acids is plotted as a function of time of incubation in Figure 1. The half-life of drug release, calculated assuming a single-exponential model for the data shown in Figure 1, is given in TABLE 2. The data in TABLE 2 are the mean and standard deviation from three experiments.

TABLE 2

15	<u>Acid</u>	<u>Percent Capture of Cytarabine</u>	<u>Half-Life in Days for Release of Cytarabine</u>	<u>Symbol in Figure 1</u>
	Hydrochloric Acid	49 ± 5	65.7 ± 4.4	◆
20	Perchloric Acid	45 ± 5	37.2 ± 8.0	▼
	Nitric Acid	44 ± 3	54.5 ± 5.7	■
	Phosphoric Acid	72 ± 1	6.5 ± 0.2	▲
	No Acid	46 ± 2	5.3 ± 0.5	●
	Formic Acid	37 ± 2	5.6 ± 0.2	◇
25	Trichloroacetic Acid	29 ± 1	5.5 ± 0.6	▽
	Acetic Acid	30 ± 2	4.8 ± 0.5	□
	Trifluoroacetic Acid	35 ± 1	3.4 ± 0.4	△
30	Sulfuric Acid	57 ± 4	1.6 ± 0.5	○

It was surprising and unexpected that the nature of the acid had a profound effect on the release rates of cytarabine in human plasma. Use of monoprotic inorganic acids, namely, hydrochloric acid, nitric acid, and perchloric acid, resulted in the slowest release rate for cytarabine. Diprotic and triprotic acids, i.e., sulfuric acid and phosphoric acid, resulted in fast release rates. The organic acids, formic acid, acetic acid, trifluoroacetic acid and trichloroacetic acid, also resulted in fast release rates.

Thus, the present disclosure provides "depot" preparations of wide application and uses in which biologically active substances are encapsulated in

- 21 -

biologically active substances are encapsulated in relatively large amounts, provide prolonged exposure or delivery at therapeutic concentrations of these substances for optimal results, and the release rate of the substance is controlled by varying the nature of the acid used in the formulation.

The present invention, therefore, is well suited and adapted to attain the ends and objects and has the advantages and features mentioned as well as others inherent therein.

While presently preferred embodiments of the invention have been given for the purpose of disclosure, changes may be made therein which are within the spirit of the invention as defined by the scope of the appended claims.

CLAIMS

1. A composition comprising a synthetic membrane vesicle comprising lipid bilayer membranes enclosing multiple non-concentric aqueous chambers containing one or more biologically active substance encapsulated therein and one or more non-hydrohalide release-rate modifying agents.
2. The composition of claim 1, wherein the release-rate modifying agents are selected from the group consisting of nitric acid, perchloric acid, formic acid, sulfuric acid, phosphoric acid, acetic acid, trichloroacetic acid, and trifluoroacetic acid, and salts or combinations thereof.
3. The composition of claim 1 wherein the release-rate modifying agent is a monoprotic inorganic acid.
4. The composition of claim 2, wherein the acids are neutralized with a proton acceptor.
5. The composition of claim 1 wherein the biologically active substance is a drug. substance.
6. The composition of claim 1 wherein the biologically active substances are selected from the group consisting of antibiotics, vaccines, antivirals, antifungals, anti-tumor drugs, proteins and glycoproteins.
7. A composition of claim 6 wherein the anti-tumor drug is cytarabine.
8. A composition of claim 1 wherein the biologically active substances are selected from the group consisting of herbicides and pesticides.
9. A targeted delivery system comprising a composition of Claim 1 with a targeting ligand attached thereto.
10. A targeted delivery system of claim 9 wherein the targeting ligand is an antibody or fragment thereof.

- 23 -

11. A targeted delivery system of claim 9 wherein the antibody is a monoclonal antibody.
12. A targeted delivery system of claim 9 wherein lipid groups are incorporated into the lipid bilayer of the synthetic membrane vesicle.
13. The composition of claim 1 wherein the synthetic membrane vesicle is anatomically targeted.
14. The composition of claim 1 wherein the synthetic membrane vesicle is mechanistically targeted.
15. The composition of claim 1 wherein the synthetic membrane vesicle is passively targeted.
16. The composition of claim 1 wherein the synthetic membrane vesicle is actively targeted.
17. The composition of claim 16 wherein the synthetic membrane vesicle is actively targeted by coupling with a moiety selected from the group consisting of a sugar, a glycolipid, and a protein.

- 24 -

18. A synthetic membrane vesicle of claim 1 produced by the method comprising:
- (a) forming a water-in-oil emulsion from two immiscible components containing at least one organic solvent, water, at least one biologically active substance, and at least one non-hydrohalide release-rate modifying agent;
 - (b) dispersing the said water-in-oil emulsion into an aqueous component to form solvent spherules; and
 - (c) removing the organic solvent from the solvent spherules to form the synthetic membrane vesicle.
19. A process for producing synthetic membrane vesicles comprising the steps of:
- (a) forming a water-in-oil emulsion from two immiscible components containing at least one organic solvent, water, at least one biologically active substance, and at least one non-hydrohalide release-rate modifying agent;
 - (b) dispersing the water-in-oil emulsion into an aqueous component to form solvent spherules; and
 - (c) removing the organic solvent from the solvent spherules to form the synthetic membrane vesicles containing aqueous droplets with the biologically active substance and the release rate modifying agent dissolved therein..
20. The process of claim 19 wherein, the concentration of the non-hydrohalide release-rate modifying agent is present in the range of about 0.1 mM to about 0.5 M.
21. The process of claim 19 wherein, an acid neutralizing agent in a concentration of from about 0.1 mM to about 0.5 M is added during step (b).

- 25 -

22. The process of claim 19 where, the organic solvent has a dissolved lipid component containing at least one amphipathic lipid with a net negative charge and at least one neutral lipid.
- 5 23. The process according to claim 22 wherein the lipid component is selected from the group consisting of a phospholipid and an admixture of phospholipids.
24. The process according to claim 23 wherein, the phospholipids are selected from the group
10 consisting of phosphatidylcholine, cardiolipin, phosphatidylethanolamine, sphingomyelin, lysophosphatidylcholine, phosphatidylserine, phosphatidylinositol, phosphatidylglycerol, and phosphatidic acid.
- 15 25. The process according to claim 24 wherein, at least one of the phospholipids has at least one net negative charge.
26. The process according to claim 24 wherein, the phospholipid is provided in admixture with
20 cholesterol.
27. The process according to claim 24 wherein, the phospholipid is provided in admixture with stearylamine.
28. The process according to claim 22 wherein, a
25 lipophilic biologically active material is provided in admixture with the lipid component.
29. The process according to claim 22 wherein, the neutral lipid is selected from the group consisting of triolein, trioctanoin, vegetable oil, lard, beef
30 fat, tocopherol, and combinations thereof.
30. The process according to claim 19 wherein, the organic solvent is selected from the group
35 consisting of ethers, hydrocarbons, halogenated hydrocarbons, halogenated ethers, esters, and combinations thereof.

- 26 -

31. The process according to claim 19 wherein, the biologically active material is hydrophilic.
32. The process according to claim 19 wherein, the emulsion is formed using a method selected from the group consisting of mechanical agitation, ultrasonic energy, and nozzle atomization.
33. The process according to claim 32 wherein, the average size of the synthetic membrane vesicles and number of the aqueous chambers therewithin are determined by the type, intensity, and duration of the emulsification method selected.
34. The process according to claim 19 wherein, the release rate modifying agent is a monoprotic inorganic acid, and aqueous component contains at least one neutralizing agent.
35. The process according to claim 34 wherein, the neutralizing agent is selected from the group consisting of free-base lysine, free base histidine, and a combination thereof.
36. The process according to claim 34 wherein, the aqueous component is an aqueous solution containing solutes selected from the group consisting of carbohydrates and amino acids.
37. The process according to claim 34 wherein, the aqueous component is an aqueous solution containing solutes selected from the group consisting of glucose, sucrose, lactose, free-base lysine, free-base histidine, and combinations thereof.
38. The process according to claim 19 wherein, the solvent spherules are formed using a method selected from the group consisting of mechanical agitation, ultrasonic energy, nozzle atomization, and combinations thereof.

- 27 -

39. The process according to claim 38 wherein, the average size of the synthetic membrane vesicle is determined by the type, intensity, and duration of the energy used.
- 5 40. The process according to claim 19 wherein, the organic solvent is removed by passing gas over the aqueous component.
- 10 41. The process of Claim 19 wherein, the biologically active substance is selected from the group consisting of antiasthmas, cardiac glycosides, antihypertensives, antiparasitics, nucleic acids and analogs, antibiotics, vaccines, antiarrhythmics, antianginas, hormones, antidiabetics, antineoplastics, immunomodulators, 15 antifungals, tranquilizers, steroids, sedatives and analgesics, vasopressors, antivirals, monoclonal antibodies, herbicides, pesticides, proteins and glycoproteins, neurotransmitters, radionuclides, radio contrasts, and combinations thereof.
- 20 42. The synthetic membrane vesicle of claim 32 or 33 wherein, the biologically active substance is selected from the group consisting of antiasthmas, cardiac glycosides, antihypertensives, 25 antiparasitics, nucleic acids and analogs, antibiotics, vaccines, antiarrhythmics, antianginas, hormones, antidiabetics, antineoplastics, immunomodulators, antifungals, tranquilizers, steroids, sedatives and analgesics, vasopressors, antivirals, monoclonal antibodies, 30 herbicides, pesticides, proteins and glycoproteins, neurotransmitters, radionuclides, radio contrasts, and combinations thereof.

- 28 -

43. A method for the treatment of a patient with a biologically active compound comprising:
administering a therapeutic amount of a therapeutic agent to a patient encapsulated in a synthetic membrane vesicle in the presence of a non-hydrohalide release-rate modifying agent effective to control the rate of release of the compound at a therapeutic level.
- 5
44. A method for the treatment of a patient with a biologically active compound comprising:
administering to the patient a synthetic membrane vesicle of claims 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10.
- 10
45. The method of claim 19 wherein, the biologically active substance is selected from the group consisting of herbicides and pesticides.
- 15

1/1




FIG. 1

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 9/127

US CL :424/450

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/450, 417

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	CANCER RESEARCH 53, April 1, 1993 (KIM) "Prolongation of Drug Exposure in Cerebrospinal Fluid by Encapsulation into DepoFoam" pp. 1596-1598, see entire document.	1, 3-7, 9-10, 12-19, 21-26, 28, 30-44
Y	US, A, 4,310,506 (BALDESCHWIELER) 12 January 1982, see columns 3 and 4 and Examples II and III.	1, 19, 27
Y	US, A, 4,224,179 (SCHNEIDER) 23 September 1980, see columns 7-9.	1, 19, 27, 29
Y	US, A, 4,752,425 (MARTIN) 21 June 1988, see entire document.	19, 32
Y	US, A, 4,920,016 (ALLEN) 24 April 1990, see columns 4-22.	1, 11, 16, 17

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be part of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*&* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

19 DECEMBER 1994

Date of mailing of the international search report

09 FEB 1995

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

THURMAN K. PAGE

Telephone No. (703) 308-2351

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US, A, 5,211,955 (LEGROS) 18 May 1993, see entire document.	1, 2, 19, 20
Y	US, A, 4,588,578 (FOUNTAIN) 13 May 1986, see columns 6, lines 8-19 and column 13, lines 60-68.	8, 45